Novel Approach to Benzimidazoles Using Fe₃O₄ Nanoparticles as a Magnetically Recoverable Catalyst

Bahador KARAMI^{1,*}, Shaghayegh NIKOSERESHT², Saeed KHODABAKHSHI¹

¹Department of Chemistry, Yasouj University, Yasouj, Iran

²Department of Chemistry, Gachsaran Branch, Islamic Azad University, Gachsaran, Iran

Abstract: Magnetically recoverable Fe₃O₄ nanoparticles have been synthesized as a catalyst for the cyclocondensation of 1,2-phenylenediamines with orthoesters under solvent-free conditions. Catalyst loadings can be as low as 1 mol% to give high yields of the corresponding benzimidazole derivative at 80 °C. This green method offers significant advantages in terms of its simplicity, low catalyst loadings, high product yields, and non-toxic nature. The Fe₃O₄ nanoparticles were characterized by X-ray diffraction, transmission electron microscopy, and Fourier transform infrared spectroscopy.

Key words: ferroferric oxide nanoparticle; 1,2-phenylenediamine; orthoester; benzimidazole

CLC number: O643

Document code: A

Received 24 August 2011. Accepted 7 November 2011.

*Corresponding author. Tel: +98-741-2223048; Fax: +98-741-2242167; E-mail: karami@mail.yu.ac.ir

English edition available online at Elsevier ScienceDirect (<http://www.sciencedirect.com/science/journal/18722067>).

The challenge in chemistry is to develop practical methods with convenient conditions and reagents, and the concept of “green chemistry” is becoming ever important in the scientific community. Green chemistry is emerging as a high-priority guiding principle for organic synthesis. Recently, the preparation and application of nanoparticles (NPs) in organic synthesis has been the subject of intense interest [1–4]. Using NPs offers advantages for “clean” chemistry, since, in addition to being readily recovered, they are non-toxic and widely accessible. The NP surface can also be functionalized to append catalytically active groups [5,6].

In this work, we describe a new and green strategy for the preparation of novel and known benzimidazoles using Fe₃O₄ nanoparticles. These have proved as powerful, safe, and recyclable catalysts under solvent-free conditions. In a previous study, Goosen and co-workers [7] reported the catalytic decarboxylative cross-ketonisation of aryl- and alkylcarboxylic acids using Fe₃O₄ nanoparticles.

In recent years, the development of improved synthetic routes to heterocyclic compounds has attracted special attention [8–10]. Among the wide variety of nitrogen heterocycles that have been explored as pharmaceutically important compounds, benzimidazoles exhibit relatively high biological activities. They have exhibited activity against cancer [11], HIV [12,13], herpes (HSV-1) [14], influenza [15], certain fungi [16], and raf kinase [17]. The most prominent benzimidazole compound in nature is *N*-ribosyldimethylbenzimidazole, which serves as an axial

ligand for cobalt in vitamin B₁₂ [18].

Until now, several methods for the synthesis of benzimidazoles and derivatives thereof have been reported. The most common method is the condensation of an aryl-enediamine with a carbonyl equivalent [19,20]. Esters, lactones, and anhydrides can also produce benzimidazoles through the cyclization of amide [21]. However, some of these reported methods suffer from drawbacks, such as the requirement for toxic and expensive catalysts and solvents, long reaction times, low yields, and difficult work-up procedures.

1 Experimental

1.1 Preparation of Fe₃O₄ nanoparticles

FeCl₃ (5.2 g) and FeCl₂ (2.0 g) were successively dissolved in 25 ml distilled water with 0.85 ml 12.1 mol/L HCl. The resulting solution was added dropwise to a 250 ml 1.5 mol/L NaOH solution with vigorous stirring. The last step generates an instant black precipitate. The precipitate was isolated in a magnetic field, and the supernatant was removed from the precipitate through decantation [22].

1.2 Characterization of Fe₃O₄ nanoparticles

Transmission electron microscopy (TEM) studies were carried out with a JEOL JEM 3010 instrument, operating at an accelerating voltage of 300 kV. X-ray diffraction (XRD,

D8, Advance, Bruker, axs) patterns were obtained for the characterization of the heterogeneous catalysts. Fourier transform infrared (FT-IR) spectra were recorded on a Shimadzu-470 spectrometer using KBr pellets.

1.3 Synthesis of benzimidazole derivatives

A mixture of 1,2-phenylenediamine **1** (1 mmol), orthoester **2** (1 mmol), and Fe₃O₄ nanoparticles (2.32 mg, 1 mol%) was stirred and heated at 80 °C for the appropriate

Scheme 1. Synthesis of benzimidazole derivatives using Fe₃O₄ nanoparticles.

All synthesized compounds (**3a–3i**) were characterized via analytical and spectroscopic methods. Melting points were determined on a KRUSS model instrument. ¹H NMR spectra were recorded on a Bruker Avance II 400 NMR spectrometer at 400 MHz, in which CDCl₃ was used as solvent and TMS as the internal standard. All products (except for novel compounds) were characterized through comparison with the reported in the literatures [23–27].

2 Results and discussion

Figure 1 shows the XRD pattern for the Fe₃O₄ nanoparticles. A number of prominent Bragg reflections, as assigned from their indices ((220), (311), (400), (422), (511), and (440)), reveal that the resultant nanoparticles are Fe₃O₄ with a spinel structure [28]. The size of the Fe₃O₄ nanoparticles was also determined from X-ray line broadening, using the Debye-Scherrer formula, which is given as $D = 0.9\lambda/\beta\cos\theta$, where D is the average crystalline size, λ is the applied

time. Upon completion (as adjudged by TLC, silica-gel 60 F₂₅₄, *n*-hexane:ethyl acetate), hot EtOH/H₂O (50/50, 10 ml) was added to the reaction mixture and the catalyst was separated through magnetic absorption using a magnet. After separating the catalyst, the solution was heated and filtered to afford the pure product **3** in the crystal form. At the end of the reaction, the separated catalyst was washed with diethyl ether, dried at 130 °C for 1 h, and reused in another cycle. The general procedure is shown in Scheme 1.

X-ray wavelength, β is the angular line width at half maximum intensity, and θ is the Bragg's angle. For the (311) reflection, the average size of the Fe₃O₄ nanoparticles was estimated to be around 13 nm.

Figure 2 shows the TEM image of the Fe₃O₄ nanoparticles. The TEM image reveals spherical Fe₃O₄ nanoparticles with an average size of 10 nm.

Fig. 2. TEM image for the Fe₃O₄ nanoparticles.

Fig. 1. XRD pattern for the Fe₃O₄ nanoparticles.

The FT-IR spectrum for the Fe₃O₄ nanoparticles is shown in Fig. 3. The absorbance bands at 584.3 cm⁻¹ are ascribed to Fe²⁺-O²⁻, and are consistent with those reported for spinel Fe₃O₄ [29,30].

In the past few years, we have been involved in a program directed towards developing simple, novel, and facile methods for the preparation of organic compounds using a range of catalysts and readily available starting materials [23–27]. In this study, we present a simple route towards novel and known benzimidazoles **3** through the reaction of 1,2-phenylenediamines **1** with orthoesters **2** using Fe₃O₄ nanoparticles as a magnetically recoverable catalyst (Scheme 1).

Fig. 3. FT-IR spectrum for the Fe_3O_4 nanoparticles.

To identify the suitable reaction conditions for the synthesis of benzimidazole derivatives **3** using these Fe_3O_4 nanoparticles, triethyl orthoformate and *o*-phenylene diamine were selected as the model reaction. As can be seen in Table 1, we found that in the absence of the catalyst, the reaction did not proceed, even at a high temperature. Through screening, we found that this reaction proceeded smoothly with 1 mol% catalyst at 80 °C under solvent-free conditions in only 12 min. Higher loadings of catalyst did not affect a marked influence on the product yield. This may be due to the coordination of excessive catalyst to the diamine.

Table 1 Synthesis of benzimidazole (**3a**) catalyzed with various loadings of Fe_3O_4 nanoparticles at 80 °C under solvent-free conditions

Catalyst amount (mol%)	Time (min)	Yield ^a (%)
—	540	50 ^b
1	12	91
3	12	90
5	15	90

^aIsolated yields. ^bNot completed.

To prove the general applicability of this method, after optimizing the reaction conditions, different orthoesters and *o*-phenylenediamines were examined. The results are summarized in Table 2.

Encouraged by the above results, we then focused our attention on 4-benzoyl-1,2-phenylenediamine, to prepare new benzimidazoles **3j**, **3k**, and **3l**. The results obtained are summarized in Table 3.

The ^1H NMR spectrum for **3l** gave a triplet for the protons of the methyl group ($\delta = 1.47$), a quartet for the protons of the methylene group ($\delta = 3.00$) and a set of multiplet signals ($\delta = 7.82\text{--}7.45$, $7.23\text{--}7.27$) that correspond to the aromatic protons. The resonance for the NH proton appeared as a broad signal at $\delta = 10.80\text{--}10.40$. The carbonyl groups in the ^{13}C NMR spectrum of **3l** appeared at $\delta = 197.62$.

Table 2 Synthesis of benzimidazole derivatives using 1 mol% Fe_3O_4 nanoparticles at 80 °C under solvent-free conditions

Product ^a	G	R ¹	R ²	Time (min)	Yield ^b (%)	Melting point (°C)	
						Found	Ref. [31]
3a	H	H	Et	12	91	172–174	170–172
3b	H	Me	Me	15	88	176–178	172–174
3c	H	Et	Et	17	93	166–168	163–165
3d	Me	H	Et	36	77	108–110	112–114
3e	Me	Me	Me	22	84	198–200	198–200
3f	Me	Et	Et	25	80	165–167	160–163
3g	NO_2	H	Et	7	91	208–210	206–208
3h	NO_2	Me	Me	10	82	225–227	225–227
3i	NO_2	Et	Et	8	82	178–180	177–179

^aIdentified by comparison with authentic samples. ^bIsolated yields.

Table 3 Synthesis of new benzimidazole derivatives using 1 mol% Fe_3O_4 nanoparticles at 80 °C under solvent-free conditions

Product	Orthoester	R	Time (min)	Yield ^a (%)	Melting point (°C)
3j	$\text{HC}(\text{OEt})_3$	H	17	85	136–138
3j	$\text{HC}(\text{OMe})_3$	H	14	82	136–138
3k	$\text{MeC}(\text{OMe})_3$	Me	10	88	168–170
3l	$\text{EtC}(\text{OEt})_3$	Et	12	90	140–142

^aIsolated yields.

To compare the catalytic activity of Fe_3O_4 powder with that of the Fe_3O_4 nanoparticles, two parallel reactions were performed. As shown in Table 4, the product yields were moderate (55%–75%) after 50–70 min with the Fe_3O_4 powder, whilst the Fe_3O_4 nanoparticles provided excellent yields (82%–91%) after 8–12 min under solvent-free conditions.

Table 4 Comparing the catalytic activities of Fe_3O_4 powder and Fe_3O_4 nanoparticles in the synthesis of benzimidazoles

Product	Fe_3O_4 powder		$\text{Fe}_3\text{O}_4\text{NPs}$	
	Yield ^a (%)	Time (min)	Yield ^a (%)	Time (min)
3a	75	50	91	12
3g	60	56	91	7
3i	55	70	82	8

^aIsolated yields.

The design and synthesis of recoverable catalysts is a highly challenging interdisciplinary field, which combines chemistry, materials science, and engineering from economic and environmental perspective. The main disadvantage for many of the reported methods is that the catalysts are destroyed in the work-up procedure and cannot be re-

covered or re-used. In this process, as outlined in Fig. 4, the recycled catalyst can be used in up to five cycles, during which there are negligible losses in the catalytic activity.

Fig. 4. Recyclability of the Fe₃O₄ nanoparticle catalyst in the synthesis of **3a** at 80 °C under solvent-free conditions (reaction time = 12–15 min).

3 Conclusions

A green methodology for the preparation of novel and known benzimidazole derivatives, using magnetically recoverable Fe₃O₄ nanoparticles under solvent-free conditions has been presented. This method not only provides high yields and purities, but it is also cheap, convenient, quick, and economically friendly.

Acknowledgments

The authors gratefully acknowledge partial support of this work by the Yasouj University, Iran.

References

- Shi H X, Zhang T Y, Wang H L, Wang X, He M. *Chin J Catal*, 2011, **32**: 46
- Bamoharram F F, Heravi M M, Ebrahimi J, Ahmadpour A, Zebarjad M. *Chin J Catal*, 2011, **32**: 782
- Dong Q, Ouyang L M, Liu J W, Xu J H. *Chin J Catal*, 2010, **31**: 1227
- Efe C, Lykakis I N, Stratakis M. *Chem Commun*, 2011, **47**: 803
- Laska U, Frost C G, Plucinski P K, Price G J. *Catal Lett*, 2008, **122**: 68
- Tundo P, Anastas P T. *Green Chemistry: Challenging Perspectives*. Oxford: Oxford Science, 1999
- Goosen L J, Mamone P, Oppel C. *Adv Synth Catal*, 2011, **353**: 57
- Genovese S, Epifano F, Marcotullio M C, Pelucchini C, Curini M. *Tetrahedron Lett*, 2011, **52**: 3474
- Curini M, Epifano F, Chimichi S, Montanari F, Nocchetti M, Rosati O. *Tetrahedron Lett*, 2005, **46**: 3497
- Epifano F, Pelucchini C, Rosati O, Genovese S, Curini M. *Catal Lett*, 2011, **141**: 844
- Huang S-T, Hsei I-J, Chen C. *Bioorg Med Chem*, 2006, **14**: 6106
- Porcari A R, Devivar R V, Kucera L S, Drach J C, Townsend L B. *J Med Chem*, 1998, **41**: 1252
- Roth T, Morningstar M L, Boyer P L, Hughes S H, Buckheitjr R W Jr, Michejda C J. *J Med Chem*, 1997, **40**: 4199
- Migawa M T, Girardet J-L, Walker II J A, Koszalka G W, Chamberlain S D, Drach J C, Townsend L B. *J Med Chem*, 1998, **41**: 1242
- Tamm I. *Science*, 1957, **126**: 1235
- Kus C, Altanlar N. *Turk J Chem*, 2003, **27**: 35
- Buchstaller H-P, Burgdorf L, Finsinger D, Stieber F, Sirrenberg C, Amendt C, Grell M, Zenke F, Krier M. *Bioorg Med Chem Lett*, 2011, **21**: 2264
- Barker H A, Smyth R D, Weissbach H, Toohey J I, Ladd J N, Volcani B E. *J Biol Chem*, 1960, **235**: 480
- Wright J B. *Chem Rev*, 1951, **48**: 397
- Preston P N. *Chem Rev*, 1974, **74**: 279
- Niknam K, Fatehi-Raviz A. *J Iran Chem Soc*, 2007, **4**: 438
- Kang Y S, Risbud S, Rabolt J F, Stroeve P. *Chem Mater*, 1996, **8**: 2209
- Kumar R S, Idhayadhulla A, Abdul Nasser A J, Selvin J. *J Serb Chem Soc*, 2011, **76**: 1
- Karami B, Khodabakhshi S, Nikrooz M. *Polycyclic Aromat Comp*, 2011, **31**: 97
- Mallakpour S E, Karami-Dezcho B, Sheikholeslami B. *Polym Int*, 1998, **45**: 98
- Heydari A, Larijani H, Emami J, Karami B. *Tetrahedron Lett*, 2000, **41**: 2471
- Damavandi J A, Karami B, Zolfigol M A. *Synlett*, 2002: 933
- Park J, An K, Hwang Y, Park J-G, Noh H-J, Kim J-Y, Park J-H, Hwang N-M, Hyeon T. *Nature Mater*, 2004, **3**: 891
- Liu Y S, Liu P, Su Z X, Li F S, Wen F S. *Appl Surf Sci*, 2008, **255**: 2020
- Wuang S C, Neoh K G, Kang E-T, Pack D W, Leckband D E. *J Mater Chem*, 2007, **17**: 3354
- Mohammadpoor-baltork I, Khosropour A R, Hojati S F. *Catal Commun*, 2007, **8**: 1865