available at www.sciencedirect.com

journal homepage: www.elsevier.com/locate/chnjc

Communication (Special Issue on Catalysis in Organic Synthesis)

Pd(0)-catalyzed benzylation of indole through η^3 -benzyl palladium intermediateZheng-Le Zhao ^{a,b}, Qing Gu ^b, Xin-Yan Wu ^{a,*}, Shu-Li You ^{a,b,#}^a Key Laboratory for Advanced Materials and Institute of Fine Chemicals, East China University of Science and Technology, Shanghai 200237, China^b State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, China

ARTICLE INFO

Article history:

Received 13 June 2014

Accepted 3 July 2014

Published 20 January 2015

Keywords:

Benzylation

Indole

Triarylmethane

 η^3 -Benzyl palladium intermediate

ABSTRACT

An efficient method has been developed for the Pd(0)-catalyzed benzylation of indoles, which occurred with exclusive regioselectivity. When this reaction was performed in the presence of Pd(PPh₃)₄, it provided access to a broad range of substituted indoles bearing diarylmethanes at their 3-position in 90%–99% yields under mild conditions.

© 2015, Dalian Institute of Chemical Physics, Chinese Academy of Sciences.

Published by Elsevier B.V. All rights reserved.

The chemistry of π -allyl palladium complexes has become one of the most successful areas of organic synthesis, as exemplified by the Tsuji–Trost reaction [1]. In contrast, reactions involving the η^3 -benzyl palladium complexes generated from benzylic derivatives are much less common because they require the dearomatization of an aryl system. Despite the difficulties associated with the preparation of η^3 -benzyl palladium complexes, these systems are considered to be useful intermediates in organic synthesis, because they can be used as special η^3 -allyl-palladium species. In this way, η^3 -benzyl palladium intermediates have been successfully substituted at their benzylic, *ortho*- and *para*-positions with a wide variety of different nucleophiles, including carbon [2–25], nitrogen [16,26–27], oxygen [16,28] and sulfur-based nucleophiles [16,29] (Fig. 1).

η^3 -Benzyl palladium intermediates can be generated by the oxidative addition reactions of a wide range of benzylic derivatives, including carbonates [3–7,28–31], phosphates [8–10],

carboxylates [11–15] and even fluorides [16], to Pd(0). Benzyl chlorides can also be treated as common precursors for the formation of η^3 -benzyl palladium species because they are readily available with various commercial suppliers and stable [17–27]. Several elegant palladium-catalyzed benzylation processes have been developed during the course of the last dec-

Fig. 1. η^3 -Benzyl palladium intermediates and possible products.

* Corresponding author. E-mail: xinyanwu@ecust.edu.cn

Corresponding author. Fax: +86-21-54925087; E-mail: slyou@sioc.ac.cn

This work was supported by the National Basic Research Program of China (973 Program, 2010CB833300) and the National Natural Science Foundation of China (21025209, 21121062, 21272253, 21332009).

DOI: 10.1016/S1872-2067(14)60191-1 | http://www.sciencedirect.com/science/journal/18722067 | Chin. J. Catal., Vol. 36, No. 1, January 2015

ade. In 2001, Yamamoto group [17] reported a facile palladium-catalyzed allylative dearomatization reaction involving the reaction of benzyl chloride with allyltributyltin at the *para*-position. Following on from this work, Bao group [19–27] reported their research towards the development of a series of reactions involving η^3 -benzyl palladium intermediates. Despite these developments, there have been very few studies reported in the literature pertaining to the reaction of electron-rich arenes as nucleophiles with η^3 -benzyl palladium intermediates [32,33]. In this study, it was envisioned that indole could be used as an effective nucleophile to attack benzylic systems and afford the corresponding triarylmethane compounds bearing an indole fragment, which could have numerous potential applications in organic synthesis [34–40]. It is noteworthy, however, that the use of indole in this way could also result in nucleophilic attack at the *ortho*- or *para*-position of the naphthyl ring, which would result in the formation of the corresponding dearomatized product through a catalytic dearomatization process [41–44]. Herein, we report the results of our recent study towards the Pd-catalyzed benzylation of indoles.

The reaction of 1-(chloro(phenyl)methyl)-naphthalene (**1a**) with indole (**2a**) was initially selected as a model reaction to allow for the optimization of the reaction conditions. Several different variables were evaluated, including the solvent, base and reaction time, in terms of their impact on the yield of the reaction, and the results are summarized in Table 1. Pleasingly, the reaction proceeded smoothly when it was conducted in CH₃CN at room temperature in the presence of 10 mol % Pd(PPh₃)₄ and 2.0 equivalents of Cs₂CO₃ to afford the triarylmethane product **3a** as a single regioisomer in 89% yield (Table 1, entry 1). Furthermore, none of the dearomatized product

Table 1

Optimization of the reaction conditions^a.

Entry	Solvent	Base	Time (h)	Yield ^b (%)
1	CH ₃ CN	Cs ₂ CO ₃	15	89
2	toluene	Cs ₂ CO ₃	3.0	91
3	1,4-dioxane	Cs ₂ CO ₃	25	65
4	Et ₂ O	Cs ₂ CO ₃	25	70
5	DCM	Cs ₂ CO ₃	4.0	97
6	THF	Cs ₂ CO ₃	2.0	99
7	THF	Li ₂ CO ₃	2.5	94
8	THF	Na ₂ CO ₃	2.5	88
9	THF	K ₂ CO ₃	2.0	80
10	THF	Et ₃ N	30	85
11	THF	DBU	20	65
12	THF	—	2.0	60
13 ^c	THF	Cs ₂ CO ₃	12	—
14 ^d	THF	Cs ₂ CO ₃	24	83
15 ^e	THF	Cs ₂ CO ₃	30	75
16 ^f	THF	Cs ₂ CO ₃	4.0	85

^a Reaction conditions: **1a** (0.5 mmol), **2a** (1.0 mmol), base (1.0 mmol) and Pd(PPh₃)₄ (0.025 mmol) in solvent (3 mL) at room temperature under an argon atmosphere. ^b Isolated yield. ^c No Pd(PPh₃)₄. ^d 1.5 equiv of **2a** and Cs₂CO₃. ^e 1.0 equiv of **2a** and Cs₂CO₃. ^f Pd(PPh₃)₄ (2.5 mol %).

resulting from the nucleophilic attack of the indole on the naphthalene ring was observed. Encouraged by this preliminary result, we proceeded to evaluate the effects of several other solvents (i.e., toluene, 1,4-dioxane, Et₂O, DCM and THF) and bases (i.e., Cs₂CO₃, Li₂CO₃, Na₂CO₃, K₂CO₃, Et₃N and DBU) on the yield of the reaction (Table 1, entries 2–11). The highest yield of all of the conditions tested was achieved using a combination of THF with Cs₂CO₃, which provided the desired product in 99% yield (Table 1, entry 6). It is noteworthy that the reaction proceeded in only 60% yield when it was conducted without base (Table 1, entry 12). Furthermore, the reaction did not give any of the desired product when it was conducted in the absence of Pd(PPh₃)₄, which suggested that this reaction was proceeding via an η^3 -benzyl palladium intermediate (Table 1, entry 13). Decreasing the amounts of catalyst and indole led to a reduction in the yield of the desired product (Table 1, entries 14–16). Based on these results, the optimized conditions were determined to be the following: 5 mol % Pd(PPh₃)₄, 2.0 equiv of Cs₂CO₃ and 2.0 equiv of indole in THF at room temperature.

With the optimized conditions in hand, we proceeded to evaluate the scope of this reaction using a variety of different substituted indoles and benzyl chloride-type compounds. The results of these experiments are summarized in Table 2. The electronic properties of the indole ring had no discernible impact on the outcome of the reaction, with indoles **2** bearing an electron-donating (i.e., 7-Me or 5-MeO) or electron-withdraw-

Table 2

Pd-catalyzed benzylation of indoles^a.

Entry	3 , R ¹ , R ² , R ³	Time (h)	Yield ^b (%)
1	3a , H, H, H	2.0	99
2	3b , H, 7-Me, H	5.0	90
3	3c , H, 5-MeO, H	2.0	91
4	3d , H, 4-Br, H	3.0	93
5	3e , H, 5-Br, H	2.0	96
6	3f , H, 6-Br, H	3.0	94
7	3g , H, 6-F, H	2.5	90
8	3h , H, 6-Cl, H	4.0	96
9	3i , H, H, Me	3.0	99
10	3j , 4- ⁱ Bu, H, H	2.0	98
11	3k , 4-Me, H, H	2.0	98
12	3l , 4-F, H, H	2.0	91
13	3m , 4-Br, H, H	2.5	94
14		3.0	98
15	3n , 3o	20	trace

^a Reaction conditions: **1** (0.5 mmol), **2** (1.0 mmol), Pd(PPh₃)₄ (0.025 mmol) and Cs₂CO₃ (1.0 mmol) in THF (3 mL) at room temperature under Ar atmosphere. ^b Isolated yield.

Fig. 2. X-ray crystal structure of **3I**.

ing group (i.e., 4-Br, 5-Br, 6-F, 6-Cl or 6-Br) proceeded smoothly under the optimized condition to give the corresponding benzylated products in 90%–99% yields (Table 2, entries 1–8). *N*-Methyl indole also reacted smoothly to give benzylated product **3I** in an excellent yield (Table 2, entry 9). Furthermore, the optimized conditions appeared to be general for benzyl chlorides **1** bearing a variety of different substituents on the phenyl ring, with the corresponding benzylated products being obtained in 91%–98% yields (Table 2, entries 10–13). Notably, 2-(chloro(phenyl)-methyl)-naphthalene was also well tolerated under the optimized conditions (Table 2, entry 14). In contrast, however, only trace amounts of the benzylated product **3o** were observed following the reaction of chlorodiphenylmethane with indole, which indicated that the η^3 -naphthyl palladium intermediate had been formed preferentially (Table 2, entry 15). The structure of compound **3I** was unambiguously determined by X-ray crystallography (Fig. 2).

The asymmetric benzylation of indole was also investigated using a chiral ligand. Unfortunately, however, the use of the (*R,R*)-DACH-phenyl Trost ligand (**L1**) or (*R*)-BINAP (**L2**) led to a racemic mixture of **3a**, albeit in excellent yield (Scheme 1).

Based on these results, we proposed a catalytic cycle for this transformation, which is shown in Scheme 2. Oxidative addition of 1-(chloro(phenyl)methyl)naphthalene (**1a**) to Pd(0) would generate the η^3 -naphthyl palladium intermediate **I**, which would be in equilibrium the intermediate **II**. Intermediate **I** would be preferentially attacked by the indole at the benzylic position to afford the benzylated product **3a**, together with the regeneration of the Pd(0) catalyst.

In summary, we have developed an efficient palladi-

Scheme 1. Attempted asymmetric benzylation of indole.

Scheme 2. Proposed catalytic cycle.

um-catalyzed method for the completely regioselective benzylation of indoles to give a variety of triarylmethanes in good yields under mild conditions.

References

- [1] Tsuji J. *Palladium Reagents and Catalysts*. Chichester: John Wiley & Sons, 2004. 431
- [2] Zhang X, Feng X J, Liu H S, Qin Y H, Dai Y. *Chin J Catal* (张新, 冯秀娟, 刘鹤松, 秦余虎, 戴耀. 催化学报), 2012, 33: 523
- [3] Kuwano R, Kondo Y, Matsuyama Y. *J Am Chem Soc*, 2003, 125: 12104
- [4] Kuwano R, Kondo Y. *Org Lett*, 2004, 6: 3545
- [5] Ueno S, Komiya S, Tanaka T, Kuwano R. *Org Lett*, 2012, 14: 338
- [6] Trost B M, Czabaniuk L C. *J Am Chem Soc*, 2010, 132: 15534
- [7] Mukai T, Hirano K, Satoh T, Miura M. *Org Lett*, 2010, 12: 1360
- [8] Legros J Y, Toffano M, Fiaud J C. *Tetrahedron: Asymmetry*, 1995, 6: 1899
- [9] Trost B M, Czabaniuk L C. *J Am Chem Soc*, 2012, 134: 5778
- [10] Trost B M, Czabaniuk L C. *Chem Eur J*, 2013, 19: 15210
- [11] Legros J Y, Primault G, Toffano M, Rivière M A, Fiaud J C. *Org Lett*, 2000, 2: 433
- [12] Yokogi M, Kuwano R. *Tetrahedron Lett*, 2007, 48: 6109
- [13] Assié M, Meddour A, Fiaud J C, Legros J Y. *Tetrahedron: Asymmetry*, 2010, 21: 1701
- [14] Fields W H, Chroma J J. *Org Lett*, 2010, 12: 316
- [15] Torregrosa R R P, Ariyaratna Y, Chattopadhyay K, Tunge J A. *J Am Chem Soc*, 2010, 132: 9280
- [16] Blessley G, Holden P, Walker M, Brown J M, Gouverneur V. *Org Lett*, 2012, 14: 2754
- [17] Bao M, Nakamura H, Yamamoto Y. *J Am Chem Soc*, 2001, 123: 759
- [18] Ariafard A, Lin Z Y. *J Am Chem Soc*, 2006, 128: 13010
- [19] Lu S R, Xu Z W, Bao M, Yamamoto Y. *Angew Chem Int Ed*, 2008, 47: 4366
- [20] Peng B, Feng X J, Zhang X, Ji L Y, Bao M. *Tetrahedron*, 2010, 66: 6013
- [21] Peng B, Feng X J, Zhang X, Zhang S, Bao M. *J Org Chem*, 2010, 75: 2619
- [22] Peng B, Zhang S, Yu X Q, Feng X J, Bao M. *Org Lett*, 2011, 13: 5402
- [23] Feng X J, Sun A L, Zhang S, Yu X Q, Bao M. *Org Lett*, 2013, 15: 108
- [24] Zhang X, Feng X J, Yu X Q, He R, Bao M. *Org Biomol Chem*, 2013, 11: 4016

Graphical Abstract

Chin. J. Catal., 2015, 36: 15–18 doi: 10.1016/S1872-2067(14)60191-1

Pd(0)-catalyzed benzylation of indole through η^3 -benzyl palladium intermediate

Zheng-Le Zhao, Qing Gu, Xin-Yan Wu*, Shu-Li You*
East China University of Science and Technology;
Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences

An efficient method has been developed for the Pd(0)-catalyzed benzylation of indoles, which occurred with exclusive regioselectivity. The reaction provided a broad range of triarylmethanes bearing indoles in 90%–99% yields under mild conditions when it was conducted in the presence of Pd(PPh₃)₄.

- [25] Feng X J, Song J L, Liu H S, Wang L G, Yu X Q, Bao M. *RSC Adv*, 2013, 3: 18985
 [26] Zhang S, Wang Y, Feng X J, Bao M. *J Am Chem Soc*, 2012, 134: 5492
 [27] Zhang S, Bao M. *Synlett*, 2012, 23: 2431
 [28] Kuwano R, Kusano H. *Org Lett*, 2008, 10: 1979
 [29] Kuwano R, Kondo Y, Shirahama T. *Org Lett*, 2005, 7: 2973
 [30] Johns A M, Tye J W, Hartwig J F. *J Am Chem Soc*, 2006, 128: 16010
 [31] Kuwano R. *Synthesis*, 2009: 1049
 [32] Zhu Y, Rawal V H. *J Am Chem Soc*, 2012, 134: 111
 [33] Lakhdar S, Westermaier M, Terrier F, Goumont R, Boubaker T, Ofial A R, Mayr H. *J Org Chem*, 2006, 71: 9088
 [34] Skurdyina D F, Tolkachev V N, Zenkina D I, Nikolaeva T G, Tolkahev O N. *Pharm Chem J*, 2008, 42: 443
 [35] Duxbury D F. *Chem Rev*, 1993, 93: 381
 [36] Shchepinov M S, Korshun V A. *Chem Soc Rev*, 2003, 32: 170
 [37] Nair V, Thomas S, Mathew S C, Abhilash K G. *Tetrahedron*, 2006, 62: 6731
 [38] He Q L, Sun F L, Zheng X J, You S L. *Synlett*, 2009: 1111
 [39] Sun F L, Zheng X J, Gu Q, He Q L, You S L. *Eur J Org Chem*, 2010: 47
 [40] Wu K, Jiang Y J, Fan Y S, Sha D, Zhang S Q. *Chem Eur J*, 2013, 19: 474
 [41] Pape A R, Kaliappan K P, Kündig E P. *Chem Rev*, 2000, 100: 2917
 [42] Roche S P, Porco J A Jr. *Angew Chem Int Ed*, 2011, 50: 4068
 [43] Zhuo C X, Zhang W, You S L. *Angew Chem Int Ed*, 2012, 51: 12662
 [44] Zhuo C X, Zheng C, You S L. *Acc Chem Res*, 2014, 47: 2558

通过 η^3 -苄基钯中间体零价钯催化的吲哚苄基化反应

赵正乐^{a,b}, 顾庆^b, 伍新燕^{a,*}, 游书力^{a,b,#}

^a华东理工大学结构可控先进功能材料及其制备教育部重点实验室和精细化工研究所, 上海200237

^b中国科学院上海有机化学研究所金属有机国家重点实验室, 上海200032

摘要: 本文报道了零价钯催化的吲哚苄基化反应, 获得了优秀的区域选择性. 使用Pd(PPh₃)₄作为催化剂, 在温和反应条件下, 以90%–99%的收率获得含有各类取代吲哚片段的三芳基甲烷化合物.

关键词: 苄基化; 吲哚; 三芳基甲烷; η^3 -苄基钯中间体

收稿日期: 2014-06-13. 接受日期: 2014-07-03. 出版日期: 2015-01-20.

*通讯联系人. 电子信箱: xinyanwu@ecust.edu.cn

#通讯联系人. 传真: (021)54925087; 电子信箱: slyou@sioc.ac.cn

基金来源: 国家重点基础研究发展计划(973计划, 2010CB833300); 国家自然科学基金(21025209, 21121062, 21272253, 21332009).

本文的英文电子版由Elsevier出版社在ScienceDirect上出版(<http://www.sciencedirect.com/science/journal/18722067>).