available at www.sciencedirect.com

journal homepage: www.elsevier.com/locate/chnjc

Article

Photocatalytic hydrogen-evolution dimerization of styrenes to synthesize 1,2-dihydro-1-arylnaphthalene derivatives using Acr^+ -Mes and cobaloxime catalysts

Wenxiao Cao ^{a,b}, Chengjuan Wu ^a, Tao Lei ^a, Xiulong Yang ^a, Bin Chen ^{a,b,*}, Chenho Tung ^{a,b}, Lizhu Wu ^{a,b,#}

^a Key Laboratory of Photochemical Conversion and Optoelectronic Materials, Technical Institute of Physics and Chemistry, Chinese Academy of Sciences, Beijing 100190, China

^b University of Chinese Academy of Sciences, Beijing 100049, China

ARTICLE INFO

Article history:

Received 20 April 2018

Accepted 27 April 2018

Published 5 July 2018

Keywords:

Hydrogen-evolution dimerization
Styrenes

1,2-Dihydro-1-arylnaphthalene
derivatives

Photocatalysis

Cobaloxime catalysts

ABSTRACT

We report a hydrogen-evolution dimerization of styrenes *via* the synergistic merger of Acr^+ -Mes photocatalyst and cobaloxime proton reduction catalysts. By utilizing this dual catalyst system, 1,2-dihydro-1-arylnaphthalene derivatives can be directly constructed from commercially available styrenes. Our reaction proceeds smoothly under mild conditions without the need for oxidants or hydrogen atom transfer reagents, and the sole byproduct is hydrogen gas. Mechanistic investigation suggests that the reaction is initiated by photoinduced electron transfer under visible-light irradiation.

© 2018, Dalian Institute of Chemical Physics, Chinese Academy of Sciences.
Published by Elsevier B.V. All rights reserved.

1. Introduction

1,2-Dihydro-1-arylnaphthalene derivatives are common structural units in many biologically active natural products and pharmaceuticals [1–6]. This skeleton contains an alkene C=C bond that can be easily derivatized; for example, podophyllotoxin with high anticancer activity can be constructed by using this skeleton as a precursor [7, 8]. Over the past decades, a variety of effective methods such as Friedel-Crafts reaction

[9], Heck reaction [10], metal-catalyzed intramolecular epoxidation [11], and others [12, 13] have been developed for the construction of 1,2-dihydro-1-arylnaphthalenes [14–21]. Nevertheless, these methods often require substrate pre-functionalization and high temperatures, and have limited product regioselectivity. In recent years, visible-light-induced chemical transformations have attracted increasing interest from organic chemists because of their green and sustainable features [22–27]. Despite their high oxidation potential, sty-

* Corresponding author. Tel/Fax: +86-10-82543576; E-mail: chenbin@mail.ipc.ac.cn

Corresponding author. Tel/Fax: +86-10-82543580; E-mail: lzwu@mail.ipc.ac.cn

This work was supported by the Ministry of Science and Technology of China (2014CB239402, 2017YFA0206903), the National Natural Science Foundation of China (21390404), the Strategic Priority Research Program of the Chinese Academy of Science (XDB17000000), the Key Research Program of Frontier Sciences, the Chinese Academy of Sciences (QYZDY-SSW-JSC029).

DOI: 10.1016/S1872-2067(18)63095-5 | http://www.sciencedirect.com/science/journal/18722067 | Chin. J. Catal., Vol. 39, No. 7, July 2018

renes can undergo dimerization by acridinium-based organic dyes in the presence of a stoichiometric oxidant or hydrogen atom transfer (HAT) reagent [28,29]. 9-Mesityl-10-acridinium-based organic dye shows a strong absorption band in the visible region ($\lambda = 430$ nm) and high oxidizing ability in the excited state, and has been successfully employed in the transformation of a diverse range of alkenes into cation radicals under irradiation [30–34]. Huang *et al.* [28] recently reported the [4+2] cycloaddition of styrenes by using 9-mesityl-10-phenylacridinium tetrafluoroborate (Ph-Acr⁺-Mes BF₄[−]) as the photosensitizer and PhSSPh as an HAT reagent for the synthesis of 1,2,3,4-tetrahydro-arylnaphthalenes (Scheme 1(a)). Liang and co-workers [29] took the lead in the oxidative synthesis of 1,2-dihydro-1-arylnaphthalene derivatives from styrenes by using 9-mesityl-10-methyl acridinium perchlorate (Acr⁺-Mes ClO₄[−]) as the photosensitizer, and 2-phenylmalononitrile (PMN) and Selectfluor as the oxidants (Scheme 1(b)). However, the use of an HAT reagent always leads to the reduced product (tetralins) rather than 1,2-dihydronaphthalenes. The use of an oxidant, on the other hand, can result in overoxidized products, naphthalene derivatives. We contemplated whether we could combine catalytic amounts of the Acr⁺-Mes photosensitizer and cobalt catalyst [35–37] to synthesize 1,2-dihydro-1-arylnaphthalene derivatives by a cross-coupling hydrogen-evolution (CCHE) reaction [38–50], which avoids the use of any sacrificial oxidant and yields hydrogen gas (H₂) as the sole byproduct. To our delight, the use of Acr⁺-Mes ClO₄[−] and cobaloxime catalyst Co(dmgH)₂pyCl helped us execute our idea under visible-light irradiation at room temperature.

2. Experimental

2.1. General methods

¹H NMR spectra were recorded using a Bruker Avance DPX 400MHz instrument with tetramethylsilane (TMS) as the internal standard. ¹³C NMR spectra were obtained at 100 MHz and referenced to the internal solvent signals. Hydrogen gas was analyzed by gas chromatography (7890-II, Tianmei, China)

Scheme 1. Representative synthesis of 1-arylnaphthalene derivatives via photocatalysis.

using a thermal conductivity detector (TCD) with a 5 Å molecular sieve column and nitrogen as the carrier gas. Mass spectra were recorded using a Trio-2000 GC-MS spectrometer. Commercially available reagents and solvents were used without further purification. Blue LEDs (3 W, $\lambda = 440 \pm 10$ nm, 145 lm @700mA) were used as the irradiation light sources.

2.2. General procedure for the preparation of 1,2-dihydro-1-arylnaphthalene

A 10 mL Pyrex tube equipped with a magnetic stir bar was charged with styrene (73 μ L, 0.7 mmol) Acr⁺-Mes (8.82 mg, 0.003 mmol), Co(dmgH)₂pyCl (19.6 mg, 0.007 mmol), NaH₂PO₄ (210 mg, 1.75 mmol), and dry MeCN (5 mL). Then, the system was degassed with a stream of argon for 30 min. The mixture was irradiated by blue LEDs under an argon atmosphere for 24 h. After removal of the solvent under vacuum, the residue was purified by silica gel column chromatography using petroleum ether/ethyl acetate as the eluent to obtain the products.

1,2-dihydro-1-arylnaphthalene (**2**): colorless oil, yield 60%, petroleum ether. ¹H NMR (400 MHz, CDCl₃) δ 7.33 (t, $J = 8.0$ Hz, 2H), 7.27–7.24 (m, 3H), 7.20 (t, $J = 7.6$ Hz, 1H), 7.13–7.08 (m, 2H), 6.83 (d, $J = 7.6$ Hz, 1H), 6.56 (d, $J = 9.6$ Hz, 1H), 6.04–6.00 (m, 1H), 4.16 (t, $J = 8.8$ Hz, 1H), 2.69–2.64 (m, 2H). ¹³C NMR (101 MHz, CDCl₃) δ 144.5, 137.9, 134.1, 128.5, 128.4, 128.0, 127.9, 127.3, 127.2, 126.8, 126.5, 126.2, 43.9, 32.0. HRMS (EI) calcd. for C₁₆H₁₄ [M]: 206.1096, found: 206.1099.

7-Methyl-1-(*p*-tolyl)-1,2-dihydronaphthalene (**3**): colorless oil, yield 66%, petroleum ether. ¹H NMR (400 MHz, CDCl₃) δ 7.04–7.00 (m, 4H), 6.94–6.87 (m, 2H), 6.57 (s, 1H), 6.41 (d, $J = 9.6$ Hz, 1H), 5.84–5.80 (m, 1H), 3.96 (t, $J = 7.6$ Hz, 1H), 2.54–2.50 (m, 1H), 2.49–2.46 (m, 1H), 2.24 (s, 3H), 2.12 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 141.6, 137.9, 137.0, 135.9, 131.5, 129.1, 128.6, 128.3, 127.8, 127.3, 126.1, 126.0, 43.5, 32.2, 21.3, 21.0. HRMS (EI) calcd. for C₁₈H₁₈ [M]: 234.1409, found: 234.1412.

5-Methyl-1-(*o*-tolyl)-1,2-dihydronaphthalene (**4**): colorless oil, yield 71%, petroleum ether. ¹H NMR (400 MHz, CDCl₃) δ 7.21 (t, $J = 3.2$ Hz, 1H), 7.16–7.13 (m, 2H), 7.11 (t, $J = 6.4$ Hz, 1H), 7.01 (d, $J = 7.6$ Hz, 1H), 6.95 (t, $J = 6.4$ Hz, 1H), 6.78 (d, $J = 9.6$ Hz, 1H), 6.54 (d, $J = 7.6$ Hz, 1H), 6.10–6.05 (m, 1H), 4.38–4.34 (m, 1H), 2.59–2.54 (m, 2H), 2.39 (s, 3H), 2.36 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 142.2, 138.1, 136.2, 133.2, 132.6, 130.4, 128.5, 128.4, 127.6, 126.9, 126.3, 126.1, 125.3, 124.7, 40.1, 30.2, 19.8, 19.3. HRMS (EI) calcd. for C₁₈H₁₈ [M]: 234.1409, found: 234.1409.

4-(2,5-Dimethylphenyl)-5,8-dimethyl-1,2-dihydronaphthalene (**5**): colorless oil, isolated yield 55%, petroleum ether. ¹H NMR (400 MHz, CDCl₃) δ 7.12 (d, $J = 15.2$ Hz, 1H), 7.08 (s, 1H), 7.03 (d, $J = 8.0$ Hz, 1H), 6.94 (d, $J = 7.8$ Hz, 1H), 6.90 (s, 1H), 6.47 (s, 1H), 5.91–5.71 (m, 1H), 4.39 (d, $J = 8.9$ Hz, 1H), 2.87 (ddd, $J = 11.4, 6.0, 3.2$ Hz, 1H), 2.48 (s, 3H), 2.44 (s, 3H), 2.41–2.37 (m, 1H), 2.14 (s, 3H), 2.00 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 141.2, 135.9, 135.1, 133.6, 132.8, 131.8, 131.0, 130.3, 129.3, 128.9, 128.4, 126.8, 125.3, 125.1, 34.6, 29.9, 21.3, 19.5, 18.9, 18.4. HRMS (EI) calcd. for C₂₀H₂₂ [M]: 262.1722, found: 262.1711.

7-Methyl-4-(*m*-tolyl)-1,2-dihydronaphthalene (**6**) and

5-methyl-4-(*m*-tolyl)-1,2-dihydro-naphthalene (**6'**): colorless oil, isolated yield 76%, petroleum ether. ^1H NMR (400 MHz, CDCl_3) δ 7.86–7.74 (m, 2H), 7.46 (m, 1H), 7.41–7.32 (m, 2H), 7.30 (m, 3H), 6.98–6.91 (m, 3H), 6.87 (m, 1H), 6.78 (m, 1H), 6.70 (m, 1H), 6.50 (t, 2H), 6.03–5.95 (m, 1H), 5.80–5.70 (m, 1H), 4.20 (t, J = 8.4 Hz, 1H), 4.05 (t, J = 8.9 Hz, 1H), 2.91 (m, 2H), 2.62–2.57 (m, 2H), 2.32 (s, 3H), 2.30 (s, 3H), 2.26 (s, 3H), 2.09 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 144.7, 137.9, 137.6, 137.0, 136.2, 136.1, 135.5, 134.1, 134.0, 130.8, 130.4, 129.6, 129.2, 128.5, 128.3, 128.0, 127.8, 127.7, 127.3, 127.2, 126.9, 126.7, 125.5, 125.1, 124.9, 124.5, 124.3, 43.4, 38.4, 32.4, 32.2, 21.6, 21.0. HRMS (EI) calcd. for $\text{C}_{18}\text{H}_{18}$ [M]: 234.1409, found: 234.1409

6-Methoxy-4-(4-methoxyphenyl)-1,2-dihydronaphthalene (**7**): colorless oil, yield 47%, petroleum ether. ^1H NMR (400 MHz, CDCl_3) δ 7.14 (m, 3H), 7.02 (s, 1H), 6.84 (d, J = 1.8 Hz, 2H), 6.70 (m, 1H), 6.39 (m, 1H), 5.94–5.73 (m, 1H), 4.14–3.97 (m, 1H), 3.78 (s, 3H), 3.69 (s, 3H), 2.27 (d, J = 3.3 Hz, 1H), 2.06 (d, J = 4.8 Hz, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 159.1, 157.9, 140.0, 137.0, 129.3, 127.5, 127.4, 127.2, 124.7, 114.5, 113.9, 113.7, 111.3, 111.0, 55.4, 54.9, 48.0, 26.0. HRMS (EI) calcd. for $\text{C}_{18}\text{H}_{18}\text{O}_2$ [M]: 266.1307, found: 266.1309

6-Ethoxy-4-(4-ethoxyphenyl)-1,2-dihydronaphthalene (**8**): colorless oil, isolated yield 41%, petroleum ether. ^1H NMR (400 MHz, CDCl_3) δ 7.13 (s, 1H), 7.11 (s, 1H), 7.01 (d, J = 8.2 Hz, 1H), 6.80 (s, 1H), 6.68 (m, 1H), 6.48 (d, J = 9.6 Hz, 1H), 6.39 (d, J = 2.4 Hz, 1H), 5.94–5.78 (m, 1H), 4.01 (m, 2H), 3.90 (m, 2H), 3.44 (m, 1H), 2.58–2.53 (m, 1H), 2.27–2.25 (m, 1H), 2.07 (t, J = 5.0 Hz, 1H), 1.39 (s, 3H), 1.32 (t, J = 7.0 Hz, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 158.2, 157.6, 157.4, 140.2, 136.4, 136.2, 129.5, 127.6, 127.0, 124.6, 115.0, 114.8, 114.3, 111.7, 63.3, 63.0, 43.5, 26.0, 15.0, 14.8. HRMS (EI) calcd. for $\text{C}_{20}\text{H}_{22}\text{O}_2$ [M]: 294.1620, found: 294.1620

6-(*tert*-Butyl)-4-[4-(*tert*-butyl)phenyl]-1,2-dihydronaphthalene (**9**): white solid, isolated yield 62%, petroleum ether. ^1H NMR (400 MHz, CDCl_3) δ 7.32 (d, J = 8.3 Hz, 2H), 7.26–7.23 (m, 1H), 7.14 (m, 2H), 7.07 (d, J = 8.0 Hz, 1H), 6.96 (s, 1H), 6.53 (d, J = 9.6 Hz, 1H), 5.99–5.91 (m, 1H), 4.12 (t, J = 7.8 Hz, 1H), 2.75–2.69 (m, 1H), 2.66–2.61 (m, 1H), 1.34 (s, 9H), 1.25 (s, 9H). ^{13}C NMR (101 MHz, CDCl_3) δ 150.2, 149.0, 141.6, 137.4, 131.6, 127.8, 127.7, 126.3, 125.7, 125.3, 125.1, 123.5, 43.3, 34.6, 34.4, 31.4, 31.3. HRMS (EI) calcd. for $\text{C}_{24}\text{H}_{30}$ [M]: 318.2348, found: 318.2350

6-Fluoro-4-(4-fluorophenyl)-1,2-dihydronaphthalene (**10**): colorless oil, yield 67%, petroleum ether/ethyl acetate = 100/1. ^1H NMR (400 MHz, CDCl_3) δ 7.19–7.15 (m, 2H), 7.06 (m, 1H), 7.04–6.98 (m, 2H), 6.88–6.84 (m, 1H), 6.52–6.48 (m, 2H), 5.98–5.93 (m, 1H), 4.10–4.07 (m, 1H), 2.64–2.58 (m, 1H), 2.57–2.49 (m, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 163.0, 160.5, 140.3, 139.3, 130.3, 129.8, 127.6, 127.1, 126.2, 115.5, 115.0, 113.5, 43.3, 31.6. HRMS (EI) calcd. for $\text{C}_{16}\text{H}_{12}\text{F}_2$ [M]: 242.0907, found: 242.0909

6-Chloro-4-(4-chlorophenyl)-1,2-dihydronaphthalene (**11**): colorless oil, isolated yield 62%, petroleum ether/ethyl acetate = 100/1. ^1H NMR (400 MHz, CDCl_3) δ 7.26 (d, J = 2.5 Hz, 2H), 7.12 (dd, J = 6.9, 4.9 Hz, 2H), 7.00 (dd, J = 10.1, 8.3 Hz, 2H), 6.80–6.74 (m, 1H), 6.49 (d, J = 9.6 Hz, 1H), 6.10–5.79 (m, 1H),

4.06–3.98 (m, 1H), 2.89–2.74 (m, 1H), 2.59–2.49 (m, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 142.1, 139.2, 132.7, 132.6, 132.5, 129.7, 129.2, 128.8, 128.7, 128.0, 127.9, 127.4, 127.3, 127.2, 127.1, 43.2, 31.6. HRMS (EI) calcd. for $\text{C}_{16}\text{H}_{12}\text{Cl}_2$ [M]: 274.0316, found: 274.0353

5-Chloro-1-(*o*-chlorophenyl)-1,2-dihydronaphthalene (**12**): colorless oil, isolated yield 50%, petroleum ether/ethyl acetate = 100/1. ^1H NMR (400 MHz, CDCl_3) δ 7.40 (dd, J = 7.4, 2.0 Hz, 1H), 7.25–7.20 (m, 2H), 7.17–7.16 (m, 1H), 7.15–7.13 (m, 1H), 7.03–7.01 (m, 1H), 7.01–6.98 (m, 1H), 6.72 (d, J = 7.4 Hz, 1H), 6.10–6.05 (m, 1H), 4.69 (t, J = 8.0 Hz, 1H), 2.68–2.63 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 140.8, 138.8, 133.7, 131.9, 131.5, 129.9, 129.7, 128.7, 128.1, 127.9, 127.8, 126.9, 126.5, 125.2, 124.0, 39.9, 29.7. HRMS (EI) calcd. for $\text{C}_{16}\text{H}_{12}\text{Cl}_2$ [M]: 274.0316, found: 274.0266

6-Bromo-4-(4-bromophenyl)-1,2-dihydronaphthalene (**13**): colorless oil, isolated yield 62%, petroleum ether/ethyl acetate = 100/1. ^1H NMR (400 MHz, CDCl_3) δ 7.43 (d, J = 8.0 Hz, 2H), 7.31 (d, J = 10.0 Hz, 1H), 7.21 (dd, J = 8.4, 8.8 Hz, 1H), 7.07 (d, J = 8.4 Hz, 2H), 6.97 (d, J = 4.0 Hz, 1H), 6.48 (d, J = 9.6 Hz, 1H), 6.02–5.99 (m, 1H), 4.08–4.03 (m, 1H), 2.63–2.60 (m, 1H), 2.57–2.53 (m, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 142.6, 139.3, 132.9, 131.7, 131.6, 130.7, 130.0, 127.6, 127.8, 127.4, 127.2, 120.6, 43.1, 31.6. HRMS (EI) calcd. for $\text{C}_{16}\text{H}_{12}\text{Br}_2$ [M]: 361.9306, found: 361.9251

5-Bromo-1-(*o*-bromophenyl)-1,2-dihydronaphthalene (**14**): colorless oil, isolated yield 48%, petroleum ether/ethyl acetate = 100/1. ^1H NMR (400 MHz, CDCl_3) δ 7.59 (dd, J = 7.9, 1.1 Hz, 1H), 7.42 (dd, J = 12.6, 4.7 Hz, 2H), 7.35–7.27 (m, 1H), 7.22–7.12 (m, 1H), 7.06 (dd, J = 7.7, 1.9 Hz, 1H), 6.97–6.90 (m, 1H), 6.75 (d, J = 7.8 Hz, 1H), 6.14–5.91 (m, 1H), 4.66 (t, J = 8.1 Hz, 1H), 2.85 (d, J = 6.2 Hz, 1H), 2.63 (m, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 144.1, 142.4, 136.4, 133.0, 131.9, 130.1, 129.4, 128.4, 127.7, 127.0, 126.9, 126.6, 126.0, 123.6, 42.4, 30.0. HRMS (EI) calcd. for $\text{C}_{16}\text{H}_{12}\text{Br}_2$ [M]: 361.9306, found: 361.9183.

6-Ethoxy-4-(4-ethoxyphenyl)-1,2-dihydronaphthalene (**15**): colorless oil, yield 53%, petroleum ether/ethyl acetate = 100/1. ^1H NMR (400 MHz, CDCl_3) δ 7.21 (d, J = 8.5 Hz, 2H), 7.15–7.03 (m, 2H), 7.01 (dd, J = 13.7, 8.4 Hz, 2H), 6.90 (dd, J = 8.1, 2.2 Hz, 1H), 6.51 (d, J = 11.4 Hz, 1H), 6.06–5.91 (m, 1H), 4.08 (dt, J = 50.1, 25.2 Hz, 1H), 2.58 (s, 1H), 2.56 (s, 1H), 2.31–2.24 (m, 3H), 2.18 (d, J = 6.8 Hz, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 169.5, 169.5, 149.6, 149.4, 141.3, 139.2, 131.9, 129.4, 127.2, 126.9, 121.6, 121.0, 119.9, 43.5, 31.8, 21.2. HRMS (EI) calcd. for $\text{C}_{18}\text{H}_{18}\text{O}_4$ [M]: 322.1205, found: 322.1205

6-Benzyl-4-(4-benzylphenyl)-1,2-dihydronaphthalene (**16**): colorless oil, isolated yield 60%, petroleum ether/ethyl acetate = 100/1. ^1H NMR (400 MHz, CDCl_3) δ 7.50 (s, 1H), 7.44 (d, J = 8.1 Hz, 3H), 7.39 (d, J = 7.6 Hz, 2H), 7.33 (d, J = 7.4 Hz, 3H), 7.25 (d, J = 8.1 Hz, 3H), 7.23–7.17 (m, 3H), 7.13–7.05 (m, 2H), 6.51 (d, J = 9.6 Hz, 1H), 6.05–5.84 (m, 1H), 4.14 (t, J = 8.2 Hz, 1H), 2.71–2.56 (m, 1H), 2.05 (s, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 143.7, 143.6, 141.0, 141.1, 140.2, 139.4, 138.2, 133.4, 129.3, 128.9, 127.8, 128.7, 127.8, 127.3, 127.2, 127.1, 127.1, 127.1, 127.0, 126.9, 126.7, 125.7, 43.8, 32.1. HRMS (EI) calcd. for $\text{C}_{28}\text{H}_{22}$ [M]: 358.1722, found: 358.1722

1-(Naphthalen-1-yl)-1,2-dihydronaphthalene (**17**): white

tron-withdrawing groups or electron-donating groups could afford the corresponding 1,2-dihydro-1-arylnaphthalenes in moderate yields under standard conditions. Meanwhile, styrenes with halo group substitution at the *ortho* or *para*-position were also smoothly converted into the 1,2-dihydro-1-arylnaphthalene products in moderate yields, which have the potential for further transformation. When a *meta*-methyl styrene derivative was employed in the reaction, two region-isomers **6** and **6'** were obtained. The presence of a methoxy group or ethyloxy group at the *para*-position of styrene afforded 1,2-dihydro-1-arylnaphthalene (**7,8**) in only 47% and 41% yields because of the formation of some unidentified byproducts. Styrene with a tertiary butyl substituent at the *para*-position also gave the desired product **9** in 62% yield. Interestingly, 2-vinyl-naphthalene also worked well for this protocol and afforded the product **17** in 58% yield. However, the electron-deficient 2-pyridylethylene failed to afford the corresponding annulation product, similar to the results observed for 4-nitrostyrene and 4-cyanogroup. A large gram-scale reaction (10 mmol **1** in 50 mL dry CH₃CN) was performed to obtain 1,2-dihydro-1-arylnaphthalene **2** in 55% yield after 48 h of irradiation.

A cross-dimerization reaction using two styrenes with different substituents on the phenyl ring (e.g., a 1:1 mixture of 4-methylstyrene and 4-fluorostyrene) was also conducted (Scheme 3). The corresponding homodimerized and cross-dimerized products were obtained as a mixture (**3**:**21**:**10** = 2.2:2.1:1.0) in 57% total isolated yield, which further broadened the scope of our substrates.

Mechanistic studies were conducted to gain insights into the aforementioned transformation. When 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) was used as the radical scavenger, no desired products were formed (Scheme 4), implying that radicals are involved in our reaction process. To shed more light on the process of the reaction, we examined the interaction between styrene **1** and the excited Acr⁺-Mes. The luminescence of the excited Acr⁺-Mes could be significantly quenched by **1** with a rate constant of $9.6 \times 10^8 \text{ (mol/L)}^{-1} \text{ s}^{-1}$. On the other hand, Co(dmgH)₂pyCl could also quench the luminescence of Acr⁺-Mes with a quenching rate constant of $1.6 \times$

Scheme 3. Cross-over reaction between 4-methylstyrene and 4-fluorostyrene.

Scheme 4. Free radical capture experiment.

$10^7 \text{ (mol/L)}^{-1} \text{ s}^{-1}$, which is two orders of magnitude smaller than that of **1**. The concentration of styrene was also much higher than that of Co(dmgH)₂pyCl. Thus, we considered that the excited Acr⁺-Mes preferentially reacted with styrene than with Co(dmgH)₂pyCl (Fig. 1(a)). Furthermore, upon irradiation, the absorption spectrum (Fig. 1(b)) of the system containing styrene, Acr⁺-Mes, and Co(dmgH)₂pyCl in degassed dry MeCN immediately exhibited absorption bands at 440–500 nm and 550–650 nm, which indicated the formation of Co^{II} and Co^I species, respectively, based on reports in the literature [49].

Based on the above results, we proposed a plausible mechanism for this reaction (Scheme 5). Photosensitizer Acr⁺-Mes is

Fig. 1. (a) Emission spectra of Acr⁺-Mes (0.03 mmol/L) in the absence and respective presence of Co(dmgH)₂pyCl (0.07 mmol/L) and styrene **1** (1.0 mmol/L, short dash line) in degassed dry CH₃CN, and Acr⁺-Mes (0.03 mmol/L) in the presence of Co(dmgH)₂pyCl (0.07 mmol/L), and styrene **1** (1.0 mmol/L) in degassed dry CH₃CN (dot line). (b) UV-vis absorption spectra of a mixture of Acr⁺-Mes (0.03 mmol/L), Co(dmgH)₂pyCl (0.07 mmol/L) and styrene **1** (1.0 mmol/L) in degassed dry MeCN before and after irradiation under blue LEDs.

Scheme 5. Possible mechanism.

irradiated by visible light to reach its excited state $\text{Acr}^+-\text{Mes}^+$ ($E_{1/2}^{\text{red}} = +2.06$ V vs. SCE) [30–34], which undergoes single-electron transfer (SET) with styrene to generate styrene radical cation $1^{+\bullet}$ and Acr^+-Mes radical anion. Electrophilic trapping of intermediate $1^{+\bullet}$ by a neutral alkene molecule gives distal 1,4-radical cation **A**, which readily undergoes Friedel–Crafts annulation to form intermediate **B** via deprotonation and aromatization. The generated Acr^+-Mes radical anion donates an electron to $\text{Co}(\text{dmgH})_2\text{pyCl}$ (Co^{III}) to produce Co^{II} species and regenerate the photosensitizer Acr^+-Mes . The carbon-centered radical **B** is further oxidized by Co^{II} to afford Co^{I} species and product **2**. Protonation of the Co^{I} species provides $\text{Co}^{\text{III}}-\text{H}$ [52,53], which can react with a proton to release H_2 eventually and regenerate the generate Co^{III} catalyst.

4. Conclusions

In summary, we have developed a hydrogen-evolution dimerization methodology for the synthesis of 1,2-dihydro-1-arylnaphthalenes by combining photocatalysis and proton reduction catalysis under visible-light irradiation. By using the dual catalyst system of photosensitizer Acr^+-Mes and cobaloxime catalyst $\text{Co}(\text{dmgH})_2\text{pyCl}$, various styrenes can

be transformed into the desired products in moderate to good yields, and H_2 is the only byproduct. As opposed to previous methods, this reaction proceeds smoothly under extremely mild conditions, and no oxidized products (naphthalene derivatives) are produced. We believe that this methodology would be significant for application to a series of organic reactions.

References

- [1] C. Canel, R. M. Moraes, F. E. Dayan, D. Ferreira, *Phytochemistry*, **2000**, 54, 115–120.
- [2] H. M. L. Davies, X. Dai, M. S. Long, *J. Am. Chem. Soc.*, **2006**, 128, 2485–2490.
- [3] L. F. Silva, Jr, F. A. Siqueira, E. C. Pedrozo, F. Y. M. Vieira, A. C. Doriguetto, *Org. Lett.*, **2007**, 9, 1433–1436.
- [4] S. O. De Silva, C. St. Denis, R. Rodrigo, *J. Chem. Soc. Chem. Comm.*, **1980**, 995–997.
- [5] V. Nair, V. Sheeba, S. B. Panicker, T.G. George, R. Rajan, L. Balagopal, M. Vairamani, S. Prabhakar, *Tetrahedron*, **2000**, 56, 2461–2467.
- [6] B. L. Yvon, P. K. Datta, T.N. Le, J. L. Charlton, *Synthesis*, **2001**, 1556–1560.
- [7] T. Linker, F. Rebien, G. Toth, A. Simon, J. Kraus, G. Bringmann, *Chem. Eur. J.*, **1998**, 4, 1944–1951.
- [8] J. C. T. Reddel, K. E. Lutz, A. B. Diagne, R. J. Thomson, *Angew. Chem. Int. Ed.*, **2014**, 53, 1395–1398.
- [9] R. R. Malladi, N. S. Li, D. Tejedor, G. W. Kabalka, *Synth. Commun.*, **2000**, 30, 3613–3616.
- [10] K. Maeda, E. J. Farrington, E. Galdard, B. D. John, J. M. Brown, *Adv. Synth. Catal.*, **2002**, 344, 104–109.
- [11] C. K. Chan, N. C. Hsueh, Y. L. Tsai, M. Y. Chang, *J. Org. Chem.*, **2017**, 82, 7077–7084.
- [12] K. R. Kopecky, M. C. Hall, *Can. J. Chem.*, **1981**, 59, 3095–3104.
- [13] M. Y. Chang, Y. C. Cheng, *Org. Lett.*, **2016**, 18, 1682–1685.
- [14] J. J. M. Lamberts, T. J. H. M. Cuppen, W. H. Laarhoven, *J. Chem. Soc. Perkin Trans.*, **1985**, 1819–1827.
- [15] J. Yun, S. L. Buchwald, *J. Org. Chem.*, **2000**, 65, 767–774.
- [16] T. Kudoh, T. Mori, M. Shirahama, M. Yamada, T. Ishikawa, S. Saito, H. Kobayashi, *J. Am. Chem. Soc.*, **2007**, 129, 4939–4947.
- [17] Z. Yu, W. Dan, H. Jie, Z. Li, *Med. Chem. Res.*, **2009**, 19, 71–76.
- [18] M. L. N. Rao, D. N. Jadhav, V. Venkatesh, *Eur. J. Org. Chem.*, **2009**, 2009, 4300–4306.

Graphical Abstract

Chin. J. Catal., 2018, 39: 1194–1201 doi: 10.1016/S1872-2067(18)63095-5

Photocatalytic hydrogen-evolution dimerization of styrenes to synthesize 1,2-dihydro-1-arylnaphthalene derivatives using Acr^+-Mes and cobaloxime catalysts

Wenxiao Cao, Chengjuan Wu, Tao Lei, Xiulong Yang, Bin Chen *, Chenho Tung, Lizhu Wu *
 Technical Institute of Physics and Chemistry,
 Chinese Academy of Sciences;
 University of Chinese Academy of Sciences

By combining Acr^+-Mes and cobaloxime catalysts, hydrogen-evolution dimerization of styrenes has been achieved without the use of oxidants or HAT reagents. This reaction proceeds smoothly under mild conditions, and the sole byproduct is hydrogen gas.

- [19] V. Gudla, R. Balamurugan, *Chem. Asian J.*, **2013**, 8, 414–428.
- [20] K. Wang, L. G. Meng, Q. Zhang, L. Wang, *Green Chem.*, **2016**, 18, 2864–2870.
- [21] Y. L. Liu, M. F. Zhang, C. H. Tung, Y. F. Wang, *Acs Catal.*, **2016**, 6, 8389–8394.
- [22] E. P. Farney, T. P. Yoon, *Angew. Chem. Int. Ed.*, **2014**, 53, 793–797.
- [23] J. Liu, Q. Liu, H. Yi, C. Qin, R. P. Bai, X. T. Qi, Y. Lan, A. W. Lei, *Angew. Chem. Int. Ed.*, **2014**, 53, 502–506.
- [24] S. M. Stevenson, M. P. Shores, E. M. Ferreira, *Angew. Chem. Int. Ed.*, **2015**, 54, 6506–6510.
- [25] C. J. Wu, W. X. Cao, T. Lei, Z. H. Li, Q. Y. Meng, X. L. Yang, B. Chen, V. Ramamurthy, C. H. Tung, L. Z. Wu, *Chem. Commun.*, **2017**, 53, 8320–8323.
- [26] P. K. Pei, F. Zhang, H. Yi, A. W. Lei, *Acta Chim. Sin.*, **2017**, 75, 15–21.
- [27] X. Y. Yu, F. Zhou, J. R. Chen, W. J. Xiao, *Acta Chim. Sin.*, **2017**, 75, 86–91.
- [28] L. F. Wang, F. J. Wu, J. A. Chen, D. A. Nicewicz, Y. Huang, *Angew. Chem. Int. Ed.*, **2017**, 56, 6896–6900.
- [29] D. L. Wei, Y. R. Li and F. S. Liang, *Adv. Synth. Catal.*, **2016**, 358, 3887–3896.
- [30] S. Fukuzumi, H. Kotani, K. Ohkubo, S. Ogo, N. V. Tkachenko, H. Lemmetyinen, *J. Am. Chem. Soc.*, **2004**, 126, 1600–1601.
- [31] H. Kotani, K. Ohkubo, S. Fukuzumi, *J. Am. Chem. Soc.*, **2004**, 126, 15999–16006.
- [32] K. Ohkubo, K. Mizushima, R. Iwata, K. Souma, N. Suzuki, S. Fukuzumi, *Chem. Commun.*, **2010**, 46, 601–603.
- [33] D. A. Nicewicz, T. Nguyen, *ACS Catal.*, **2014**, 4, 355–360.
- [34] N. A. Romero, D. A. Nicewicz, *Chem. Rev.*, **2016**, 116, 10075–10166.
- [35] J. L. Dempsey, B. S. Brunschwig, J. R. Winkler, H. B. Gray, *Acc. Chem. Res.*, **2009**, 42, 1995–2004.
- [36] A. Bhattacharjee, M. Chavarot-Kerlidou, E. S. Andreiadis, M. Fontecave, M. J. Field, V. Artero, *Inorg. Chem.*, **2012**, 51, 7087–7093.
- [37] P. W. Du, R. Eisenberg, *Energy Environ. Sci.*, **2012**, 5, 6012–6021.
- [38] Q. Y. Meng, J. J. Zhong, Q. Liu, X. W. Gao, H. H. Zhang, T. Lei, Z. J. Li, K. Feng, B. Chen, C. H. Tung, L. Z. Wu, *J. Am. Chem. Soc.*, **2013**, 135, 19052–19055.
- [39] J. J. Zhong, Q. Y. Meng, B. Liu, X. B. Li, X. W. Gao, T. Lei, C. J. Wu, Z. J. Li, C. H. Tung, L. Z. Wu, *Org. Lett.*, **2014**, 16, 1988–1991.
- [40] X. W. Gao, Q. Y. Meng, J. X. Li, J. J. Zhong, T. Lei, X. B. Li, C. H. Tung, L. Z. Wu, *ACS Catal.*, **2015**, 5, 2391–2396.
- [41] M. Xiang, Q. Y. Meng, J. X. Li, Y. W. Zheng, C. Ye, Z. J. Li, B. Chen, C. H. Tung, L. Z. Wu, *Chem. Eur. J.*, **2015**, 21, 18080–18084.
- [42] G. T. Zhang, C. Liu, H. Yi, Q. Y. Meng, C. L. Bian, H. Chen, J. X. Jian, L. Z. Wu, A. W. Lei, *J. Am. Chem. Soc.*, **2015**, 137, 9273–9280.
- [43] C. J. Wu, Q. Y. Meng, T. Lei, J. J. Zhong, W. Q. Liu, L. M. Zhao, Z. J. Li, B. Chen, C. H. Tung, L. Z. Wu, *ACS Catal.*, **2016**, 6, 4635–4639.
- [44] Y. W. Zheng, B. Chen, P. Ye, K. Feng, W. G. Wang, Q. Y. Meng, L. Z. Wu, C. H. Tung, *J. Am. Chem. Soc.*, **2016**, 138, 10080–10083.
- [45] G. T. Zhang, X. Hu, C. W. Chiang, H. Yi, P. K. Pei, A. K. Singh, A. W. Lei, *J. Am. Chem. Soc.*, **2016**, 138, 12037–12040.
- [46] G. T. Zhang, L. L. Zhang, H. Yi, Y. Luo, X. T. Qi, C. H. Tung, L. Z. Wu, A. W. Lei, *Chem. Commun.*, **2016**, 52, 10407–10410.
- [47] L. B. Niu, H. Yi, S. C. Wang, T. Y. Liu, J. M. Liu, A. W. Lei, *Nat. Commun.*, **2017**, 8, 14226–16223.
- [48] H. Yi, L. B. Niu, C. L. Song, Y. Y. Li, B. W. Dou, A. K. Singh, A. W. Lei, *Angew. Chem. Int. Ed.*, **2017**, 56, 1120–1124.
- [49] X. Hu, G. T. Zhang, F. X. Bu, X. Luo, K. B. Yi, H. Zhang, A. W. Lei, *Chem. Sci.*, **2018**, 9, 1521–1526.
- [50] X. Hu, G. T. Zhang, F. X. Bu, A. W. Lei, *Angew. Chem. Int. Ed.*, **2018**, 57, 1286–1290.
- [51] P. W. Du, K. Knowles, R. Eisenberg, *J. Am. Chem. Soc.*, **2008**, 130, 12576–12577.
- [52] N. A. Romero, D. A. Nicewicz, *J. Am. Chem. Soc.*, **2014**, 136, 17024–17035.
- [53] P. A. Jacques, V. Artero, J. Pecaut, M. Fontecave, *Proc. Natl. Acad. Sci. USA*, **2009**, 106, 20627–20632.

吡啶和钴酞配合物协同光催化苯乙烯放氢二聚反应构筑 1,2-二氢-1-芳基萘

曹文晓^{a,b}, 吴成娟^a, 雷 涛^a, 杨修龙^a, 陈 彬^{a,b,*}, 佟振合^{a,b}, 吴骊珠^{a,b,#}

^a中国科学院理化技术研究所, 光化学转换与功能材料重点实验室, 北京 100190

^b中国科学院大学, 北京 100049

摘要: 芳基二氢萘类衍生物是许多生物活性的天然产物以及药物的常见结构单元, 其合成一直都受到化学家们的关注. 传统的 1,2-二氢-1-芳基萘骨架化合物的构筑大都需要进行底物的预官能团化, 在高温条件下进行, 且产物的选择性较差, 因此发展一种简单温和的制备方法很有必要. 最近兴起的可见光催化因具有条件温和、环境友好等特点而成为了合成化学家的研究热点. 近期研究发现, 在可见光作用下利用吡啶光敏剂的强氧化能力, 可以实现苯乙烯的加成. 但此类反应需要当量的氧化剂或氢原子转移试剂, 容易导致苯乙烯的二聚环合产物的进一步氧化或还原. 我们在前期发展的“放氢交叉偶联”反应的基础上, 利用吡啶光催化和钴酞催化的协同作用, 实现了苯乙烯的放氢二聚反应, 在室温下高效构筑了 1,2-二氢-1-芳基萘骨架, 反应条件温和, 底物脱除的电子和质子在钴酞催化剂作用下以氢气的形式释放, 反应具有中等及以上的收率.

本文以苯乙烯为模型底物, 吡啶为光敏剂, 钴酞配合物为质子还原催化剂, 在乙腈溶剂中, 蓝色 LED 灯下光照 24 h 可以获得 56% 的产率, 对于其它的光敏剂如 *fac*-Ir(ppy)₃ 等则不能催化该反应. 通过催化剂种类及用量筛选表明, 7 mol% 的 Co(dmgH₂)pyCl 配合物具有最好的反应效果, 可以获得 72% 的收率. 控制实验表明, 光敏剂、钴酞催化剂和光照都是必须的. 通过底物拓展我们发现, 烷基、卤素等不同取代基的苯乙烯类化合物均可以获得较好的收率, 不同苯乙烯之间也可以发生交叉反应.

随后, 我们进一步通过光谱和中间体捕获实验对反应机理进行了研究. 自由基捕获实验说明反应过程可能涉及自由基历程; 光谱淬灭实验表明苯乙烯和 Co(dmgH₂)pyCl 均可淬灭吡啶的发光, 但苯乙烯淬灭吡啶的程度远大于

Co(dmgh₂)pyCl 淬灭吡啶的程度。在反应时苯乙烯的浓度远大于催化剂的溶度, 因此, 我们认为激发态吡啶首先与苯乙烯发生反应; 可见光照射反应体系 1 min 后在 440–500 和 550–650 nm 处观察到明显的 Co^{II} 和 Co^I 的吸收峰。基于以上实验结果, 我们提出了可能的催化循环: 吡啶受光激发到达激发态后, 首先与底物苯乙烯发生单电子转移生成苯乙烯正离子自由基和吡啶阴离子自由基 Acr[•]-Mes, Acr[•]-Mes 还原 Co(dmgh₂)pyCl 生成 Co^{II} 中间体, 从而回到基态完成光催化循环。苯乙烯正离子自由基与另一分子苯乙烯加成环合, 进而通过芳构化生成自由基中间体, 再与 Co^{II} 作用生成目标产物 1,2-二氢-1-芳基萘和 Co^I, Co^I 通过结合体系中的质子进而释放出氢气回到 Co^{III} 从而完成钴卟催化循环。

关键词: 放氢二聚; 苯乙烯; 1,2-二氢-1-芳基萘; 光催化; 钴卟催化剂

收稿日期: 2018-04-20. 接受日期: 2018-04-27. 出版日期: 2018-07-05.

*通讯联系人. 电话/传真: (010)82543576; 电子信箱: chenbin@mail.ipc.ac.cn

#通讯联系人. 电话/传真: (010)82543580; 电子信箱: lzwu@mail.ipc.ac.cn

基金来源: 中国科技部 (2014CB239402, 2017YFA0206903); 国家自然科学基金 (21390404); 中科院战略性先导科技专项 (B类) 项目 (XDB17000000); 中科院前沿科学重点研究计划项目 (QYZDY-SSW-JSC029).

本文的电子版全文由Elsevier出版社在ScienceDirect上出版(<http://www.sciencedirect.com/science/journal/18722067>).